

- _____ 10. A person is about five times more likely to contract HIV through anal intercourse than through vaginal intercourse.
A True B False
- _____ 11. How many times more likely is a woman to contract HIV through vaginal intercourse if the man is infected than in the reverse situation?
A 5 times C 10 times
B 8 times D 20 times
- _____ 12. The level of HIV circulating in the blood is called 'viral load'.
A True B False
- _____ 13. What is the name of the test used to detect the presence of HIV antibodies.
A LISA C CDC
B ELISA D Western Blot
- _____ 14. The 'window period' for detecting the HIV virus is:
A 2 to 4 weeks C 5 to 20 weeks
B 6 to 12 weeks D None of the above
- _____ 15. Flu-like symptoms are reported by roughly half of those who contract HIV and generally occurs between 2 and 6 weeks after infection.
A True B False
- _____ 16. HIV-infected persons with a CD4+T cell count of _____ are classified as persons with AIDS.
A 200 C 600
B 400 D 800
- _____ 17. **Changes in the Epidemiology of HIV/AIDS Since 1995**
HIV prevalence is higher among this population than in other ethnic groups:
A Asian C African American
B Hispanic D Native American
- _____ 18. HIV/AIDS is not commonly found among heterosexuals.
A True B False
- _____ 19. In 1998, this region of the United States had the highest incidences of AIDS cases:
A Northeast C Midwest
B South D West
- _____ 20. As of June 1999, more than half of all cumulative male adult and adolescent AIDS cases were among MSMs.
A True B False

